



AAHHHHH!

**Is Terrestrial
Methane Flux
Interesting?**

Ankur R Desai
University of Wisconsin

<http://purefixion.com/attention/2006/03/cow-farts.html>


What do I (we) do?

<http://flux.aos.wisc.edu>


- Probe spatial heterogeneity in biologically-mediated surface-atmosphere exchanges from sites to regions (meters-1000s km)
 - Forests, wetlands, lakes, urban (temperate-boreal-tropical-Mediterranean-alpine, terrestrial-aquatic, management gradients)
 - **Multiple greenhouse gases (methane), esp. with eddy covariance**
 - Feedbacks from energy balance and a land surface variability on the atmospheric boundary layer and synoptic-PBL interactions in observations and models (LES, PBL, mesoscale, climate)
 - Up/down scaling across multiple measurements: eddy covariance, biometric, airborne budgets, inverse modeling, hyperspectral remote sensing (leaf to satellite)
 - Informing ecosystem and atmospheric models with diverse measurements across space (data assimilation, model informatics)
 - <http://pecanproject.org>





- Also Site PI for
 - US-PFa Park Falls Very Tall Tower (Mixed Forest/Wetland)
 - US-WCr Willow Creek Mature Hardwood
 - US-Syv Sylvania Old-growth
 - US-Los Shrub wetland
 - Lake Mendota eutropic lake tower




Methane
1.8 ppm
Atmos lifetime ~10 yrs




Carbon Dioxide
400 ppm
30-100+ years







Mauna Loa, Hawaii, United States (MLO)




Source: NOAA ESRL

Year

Graph created ESRL/GMD - 2014-April-26 04:40 am




Methane SCIAMACHY/ENVISAT 2003-2005



CH₄ column-averaged mole fraction [ppb]




[http://www.iup.uni-bremen.de/sciamachy/
NIR_NADIR_WFM_DOAS/wfmd_image_gallery_ch4.html](http://www.iup.uni-bremen.de/sciamachy/NIR_NADIR_WFM_DOAS/wfmd_image_gallery_ch4.html)




<http://media-1.web.britannica.com/eb-media/75/135075-004-105F7745.jpg>

Global Anthropogenic CH₄ Budget by Source in 2000



Total CH₄ emissions in 2000 = 282.6 Tg CH₄






Source: Picarro, Inc. / Nathan Phillips, BU

METHANE
(Fluxes = MTC/yr)




Climate sensitivity
0.3°C/doubling CH₄
18% greenhouse






Bloom et al., Science, 2010



Spahni et al. (2011) Biogeosciences






Freshwater Methane Emissions Offset the Continental Carbon Sink


David Bastviken,^{1*} Lars J. Tranvik,² John A. Downing,³ Patrick M. Crill,⁴ Alex Enrich-Prast⁵



Methane fluxes show consistent temperature dependence across microbial to ecosystem scales

Gabriel Yvon-Durocher¹, Andrew P. Allen², David Bastviken³, Ralf Conrad⁴, Cristian Gudasz^{5,6†}, Annick St-Pierre⁷, Nguyen Thanh-Duc⁸ & Paul A. del Giorgio⁷






C. Phillips, OSU

Micrometeorological measurements of methane flux in a Minnesota peatland during two growing seasons




N.J. SHURPALI^{1,2} & S.B. VERMA^{1,*}



Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique

2007


By JANNE RINNE^{1*}, TERHI RIUTTA², MARI PIHLATIE¹, MIKA AURELA³,
SAMI HAAPANALA¹, JUHA-PEKKA TUOVINEN³, EVA STINA TUUTHI²



Comparing laser-based open- and closed-path gas analyzers to measure methane fluxes using the eddy covariance method

Matteo Detto^{a,*}, Joseph Verfaillie^a, Frank Anderson^b, Liukang Xu^c, Dennis Baldocchi^a


2011, AgForMet




Testing the applicability of neural networks as a gap-filling method using CH₄ flux data from high latitude wetlands

Biogeosci, 2013

S. Dengel¹, D. Zona^{2,3}, T. Sachs⁴, M. Aurela⁵, M. Jammet⁶, F. J. W. Parmentier⁷, W. Oechel³, and T. Vesala¹





Shoemaker et al., 2013, GRL


So what do we get from a very tall CH₄ flux tower?

- Desai, A.R., Xu, K., Tian, H., Weishampel, P., Thom, J., Baumann, D., Andrews, A.E., Cook, B.D., King, J.Y., and Kolka, R., 2014. Landscape-level terrestrial methane flux observed from a very tall tower. *Agric. Forest Meteorol.*, submitted.


Tall towers offer novel approach to estimating regional fluxes




Credit: M. Rydzik



Source: B. Cook






Long-term continuous CH₄ eddy covariance is now feasible



Picarro G1301-f 122 m
CH₄/CO₂ (H₂O)


Credit: M. Rydzik



396 m
122 m
30 m
CO₂/H₂O flux

Not shown: Los Gatos for CH₄ profile/storage flux
LI-7000 (NOAA) for CO₂ profile/storage


CH₄ random uncertainty can be large but a reasonable level of detection is possible




Based on approach of Salesky et al (2012) BLM

Storage flux is more complicated for CH_4 than CO_2 NEE


CH₄ concentration black=30 blue=122 red=396




CO₂ concentration




CH₄ flux black=NEE red=flux blue=storage




CO₂ flux








Driving factors are trickier for CH₄! Temperature at daily scale...



Moisture at annual scale







$$N_* = \frac{N}{\sum_{t=N/2}^N \left[\left(1 - \frac{t}{N} \right) \rho_t^x \rho_t^y \right]}$$

Bretherton et al., 1999, J Clim; Desai, 2014, Photosyn. Res.



Chamber CH₄ fluxes show high inter and intra site variability, and scaled fluxes are ~1/3 of tower, while tower is less than a profile similarity approach




Models get seasonal pattern but not interannual variability or large emissions




Flux footprints vary in space and time over very spatially heterogeneous sources





Environmental response Functions (ERF)


Flux grids



Before:




After:




Based on: Metzger et al., 2013, Biogeosci.

ERF model shows vegetation fraction and water vapor explain CH₄ flux variations



Gridded ERF functions show significant spatial variability in CH₄ flux



So is methane interesting?

- NO: short-lifetime, small flux in most forests, only ecologically relevant for wetlands/agriculture/tropics and arctic, anthropogenic source more important, hard/expensive to measure flux well,
- YES: high short-term (policy-relevant) radiative forcing, ecosystem climate sensitivities involve CH₄ and CO₂ flux tradeoffs, tracer of microbial ecology, data and models show lots of uncertainty and invalidity of prior assumptions of fixed ratios, ...
- *What do you think?*

Thank you!

- NSF CAREER DEB #0845166
- DOE Ameriflux Network Management Program
- NEON, Inc. Service Agreement to U Wisconsin
- WLEF/ Park Falls (US-PFa) tall tower research partners: NOAA ESRL (A. Andrews, J. Kofler), USFS NRS (M. Kubiske, D. Baumann), Penn State (K. Davis), Cal Tech (P. Wennberg), COSMOS (M. Zreda), NASA GSFC (B. Cook), WI ECB (J. Ayers), Ameriflux, NEON (S. Metzger)
- Desai lab at UW: J Thom, K Xu, and others
 - <http://flux.aos.wisc.edu>
 - desai@aos.wisc.edu
 - 608-218-4208

